

This is the peer-reviewed, manuscript version of the following article:

Taylor-Brown, F. E., Cardy, T. J. A., Liebel, F. X., Garosi, L., Kenny, P. J., Volk, H. A. and De Decker, S. (2015) 'Risk factors for early post-operative neurological deterioration in dogs undergoing a cervical dorsal laminectomy or hemilaminectomy: 100 cases (2002–2014)', *The Veterinary Journal*, 206(3), 327-331.

The final version is available online via <http://dx.doi.org/10.1016/j.tvjl.2015.10.010>.

© 2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Risk factors for early post-operative neurological deterioration in dogs undergoing a cervical dorsal laminectomy or hemilaminectomy: 100 cases (2002–2014)

AUTHORS: Taylor-Brown, F. E., Cardy, T. J. A., Liebel, F. X., Garosi, L., Kenny, P. J., Volk, H. A. and De Decker, S.

JOURNAL TITLE: The Veterinary Journal

VOLUME/EDITION: 206/3

PUBLISHER: Elsevier

PUBLICATION DATE: 22 October 2015 (online)

DOI: 10.1016/j.tvjl.2015.10.010

1 **Risk factors for early postoperative neurological deterioration in dogs undergoing a**  
2 **cervical dorsal laminectomy or hemilaminectomy: 100 cases (2002-2014)**

3  
4 F.E. Taylor-Brown <sup>a\*</sup>, T.J.A. Cardy <sup>a</sup>, F.X. Liebel <sup>b</sup>, L. Garosi <sup>b</sup>, P.J. Kenny <sup>a</sup>, H.A. Volk <sup>a</sup>, S.  
5 De Decker <sup>a</sup>

6  
7 <sup>a</sup> *Clinical Science and Services, The Royal Veterinary College, Hawkshead Lane, North*  
8 *Mymms, Hatfield, Hertfordshire, AL9 7TA, UK*

9 <sup>b</sup> *Davies Veterinary Specialists, Manor Farm Business Park, Higham Gobion, SG53HR*

10  
11 \* Corresponding author. Tel: +44(0)1707 666366.

12 *E-mail address: ftaylor@rvc.ac.uk (F.E. Taylor-Brown)*

13

14 **Abstract**

15 Early postoperative neurological deterioration is a well-known complication  
16 following dorsal cervical laminectomies and hemilaminectomies in dogs. This study aimed to  
17 evaluate potential risk factors for early postoperative neurological deterioration following  
18 these surgical procedures.

19

20 Medical records of 100 dogs that had undergone a cervical dorsal laminectomy or  
21 hemilaminectomy between 2002 and 2014 were assessed retrospectively. Assessed variables  
22 included signalment, bodyweight, duration of clinical signs, neurological status before  
23 surgery, diagnosis, surgical site, extent of surgery, and duration of procedure. Outcome  
24 measures were neurological status immediately following surgery and duration of  
25 hospitalisation. Univariate statistical analysis was performed to identify variables to be  
26 included in a multivariate model.

27

28 Diagnoses included osseous associated cervical spondylomyelopathy (OACSM;  
29  $n=41$ ), acute intervertebral disk extrusion (IVDE; 31), meningioma (11), spinal arachnoid  
30 diverticulum (10) and vertebral arch anomalies (7). Overall 55% (95% CI 45.25-64.75) of  
31 dogs were neurologically worse 48 h postoperatively. Multivariate statistical analysis  
32 identified four factors significantly related to early postoperative neurological outcome.  
33 Diagnoses of OACSM or meningioma were considered the strongest variables to predict  
34 early postoperative neurological deterioration, followed by higher (more severely affected)  
35 neurological grade before surgery and longer surgery time.

36

37 This information can aid in the management of expectations of clinical staff and  
38 owners with dogs undergoing these surgical procedures.

39

40 *Keywords:* cervical; dog; dorsal; hemilaminectomy; laminectomy

## 41 **Introduction**

42 Dorsal decompressive surgery for the treatment of cervical spinal diseases is a well-  
43 established surgical approach (Sharp and Wheeler, 2005). Cervical dorsal laminectomies and  
44 hemilaminectomies are indicated for lesions located dorsally or laterally in the cervical  
45 vertebral canal. Disorders commonly treated by this approach include osseous associated  
46 cervical spondylomyelopathy (OACSM), intervertebral disc extrusions with lateralised or  
47 dorsally located disc material, vertebral arch anomalies, spinal arachnoid diverticula (SAD),  
48 or where multiple ventral intervertebral disc protrusions are present (Gill et al., 1996; de  
49 Risio et al., 2002; da Costa, 2010; De Decker et al., 2012a).

50

51 There have been differing reports pertaining to the outcome of dorsal cervical  
52 decompressive surgeries. Several reports have suggested that cervical dorsal laminectomy  
53 should be considered an invasive procedure with a high risk of postoperative morbidity, many  
54 complications and prolonged hospitalisation and recovery times (de Risio et al., 2002, da  
55 Costa, 2010, Delamide Gasper et al., 2014). The most important complication associated with  
56 this surgical approach is early postoperative neurological deterioration with prolonged  
57 recumbency after surgery. Other complications include marked tissue disruption,  
58 intraoperative haemorrhage, prolonged surgery times, excessive scar tissue formation and  
59 cardiorespiratory compromise (de Risio et al., 2002; Delamide Gasper et al., 2014). However  
60 other studies have reported more favourable outcomes with limited hospitalisation and  
61 recovery times (Gill et al., 1996; Faissler, 2011, De Decker et al., 2012a).

62

63           It remains unclear why some patients undergoing dorsal cervical procedures  
64 experience early postoperative neurological deterioration whilst others do not. The aim of this  
65 study was to evaluate potential risk factors for early neurological deterioration in dogs  
66 following cervical dorsal laminectomy or hemilaminectomy. It was hypothesised that the  
67 specific diagnosis would influence outcome, with those dogs diagnosed with acute  
68 intervertebral disc extrusions (IVDE) less likely to have early postoperative neurological  
69 deterioration following surgery compared to dogs diagnosed with the more chronic condition  
70 OACSM. Other assessed variables thought to impact on early postoperative outcome  
71 included signalment, duration of clinical signs and neurological status before surgery,  
72 surgical site, extent of surgery and duration of procedure.

73

## 74 **Materials and methods**

### 75 **Criteria for inclusion**

76           Medical records of dogs that had undergone a cervical dorsal laminectomy or  
77 hemilaminectomy at the Royal Veterinary College between 2002 and 2014 and Davies  
78 Veterinary Specialists between 2008 and 2014 were reviewed. In order to be considered for  
79 inclusion in the study each dog needed to have complete medical records and imaging studies  
80 available for review and have clear data relating to their neurological status before and after  
81 surgery. If the neurological status could not be clearly determined then cases were excluded.  
82 Further information recorded included signalment, bodyweight, duration of clinical signs  
83 prior to surgery (acute <48 h, subacute 2-7 days and chronic >7 days), diagnosis, type,  
84 location, extent (number of vertebrae operated on) and duration of the surgery, presence of  
85 perioperative complications during the period of hospitalisation and hospitalisation time.

86

### 87 **Neurological grading**

88           A scoring system modified from de Risio et al. (2002) was used to objectively grade  
89 the dogs' neurological status; normal neurological status (grade 0), cervical hyperaesthesia  
90 without neurological deficits (grade 1), mild ataxia without paresis and slight delay in  
91 postural reactions, with or without thoracic limb deficits and/or cervical hyperaesthesia  
92 (grade 2), noticeable ataxia and paresis with delayed postural reactions, with or without  
93 thoracic limb deficits and or cervical hyperaesthesia (grade 3), paresis or absent postural  
94 reactions, with or without thoracic limb deficits and or cervical hyperaesthesia, dogs are able  
95 to rise and make a few steps with assistance (grade 4), non-ambulatory tetraparesis, with or  
96 without cervical hyperaesthesia, patients are not able to rise independently (grade 5) and  
97 tetraplegia with respiratory compromise (grade 6). Deterioration in neurological status by one  
98 or more grades was defined as postoperative neurological deterioration.

99

#### 100 Diagnostic imaging

101           Included dogs underwent diagnostics including myelography, computed tomography  
102 (CT), computed tomography-myelography (CT-m) or magnetic resonance imaging (MRI)  
103 under general anaesthesia. Although general anaesthesia protocols could vary between  
104 individual cases, a commonly used protocol included premedication with a combination of  
105 acepromazine (0.01 mg/kg IV) and methadone (0.1- 0.2 mg/kg IV), followed by induction  
106 with propofol, (4 – 6 mg/kg IV) and maintenance of general anaesthesia with isoflurane in  
107 oxygen. Myelography was performed by intrathecal injection of iohexol (Omnipaque, GE  
108 Healthcare) contrast medium between the L5-L6 articulation (0.2 ml/kg with a maximal dose  
109 of 10 ml). CT imaging was performed using a 16-slice helical CT scanner (Mx8000 IDT,  
110 Philips). After completion of the transverse CT study, sagittal and dorsal reconstructions were  
111 made. MRI was performed with a 1.5 T (Intera, Philips Medical Systems) or 0.4 T (Aperto

112 MRI, Hitachi) and included a minimum of T2- and T1-weighted sagittal and transverse  
113 images.

114

115 Localisation and categorisation of spinal cord compression

116 Information obtained from both the radiology and surgery reports was used in order to  
117 determine the site of spinal cord compression and confirm the diagnosis. In addition a board-  
118 certified neurologist (SDD) reviewed the imaging studies for diagnostic accuracy. Dogs were  
119 divided into four categories based on their diagnosis: 1: OACSM, 2: acute IVDE, 3:  
120 histopathologically confirmed meningioma, 4: SAD, 5: vertebral arch anomalies. Vertebral  
121 arch anomalies were defined as a well-defined and smooth hypertrophy of the dorsal lamina  
122 and spinous process of  $\geq 2$  adjacent vertebrae. No other osseous abnormalities were present  
123 in these dogs (De Decker et al., 2012). Lesions were classified according to their location  
124 within the cervical vertebral column with cranial lesions classified as those located between  
125 C1 and C4 vertebrae and caudal lesions classified as those from C4 to T1 vertebrae. If lesions  
126 affected both the cranial and caudal cervical vertebral column this was documented.

127

128 Surgery, postoperative care and outcome measures

129 All dogs had a dorsal laminectomy, hemilaminectomy or combination. The  
130 procedures were carried out by a board certified neurologist using published techniques  
131 (Sharp and Wheeler, 2005; Platt and da Costa, 2012). Anaesthesia protocols varied for  
132 individual dogs based on attending anaesthetist preference and specific patient requirements  
133 but typically included acepromazine (0.01 mg/kg IV) and methadone (0.1- 0.2 mg/kg IV),  
134 followed by induction with propofol, (4 – 6 mg/kg IV) and maintenance of general  
135 anaesthesia with isoflurane in oxygen. Perioperative analgesia included a combination of  
136 opioids, ketamine and non-steroidal anti-inflammatory drugs (NSAID). Postoperative

137 analgesia typically included opioids and NSAID. Most patients received postoperative  
138 physiotherapy. All dogs underwent a daily neurological assessment by a board certified  
139 neurologist and information pertaining to their assessment was recorded in the medical  
140 records.

141

#### 142 Statistical analysis

143 All variables were treated as categorical except for age, weight, number of vertebrae  
144 on which surgery was performed, duration of surgery and duration of hospitalisation, which  
145 were continuous. Early postoperative neurological deterioration by one or more grades was  
146 defined as the primary outcome measure. A secondary outcome measure was defined as  
147 duration of hospitalisation. Univariate analysis identified variables associated with early  
148 postoperative neurological deterioration. Statistical comparisons between mean values of  
149 normally distributed data were made using a one-way analysis of variance (ANOVA), with  
150 additional pairwise comparisons with Bonferroni adjustment as required for significant  
151 variables. Median values for non-parametric data were compared with either Mann-Whitney  
152 or Kruskal-Wallis tests with post-tests as required. Statistically significant results are  
153 displayed where  $P < 0.05$ . Unless otherwise stated normally distributed data is presented as  
154 mean  $\pm$  standard deviation and non-parametric data as median and range. Computations were  
155 performed using SPSS (Statistical Package for the Social Sciences v. 21.0.1; SPSS Inc.).

156

157 Prior to inclusion in a binary logistic regression model multinomial modelling was  
158 performed to identify associations between input variables. All biologically important  
159 confounders (age, weight, surgery time) and independent factors (onset of clinical signs,  
160 neurological grade preoperatively) with  $P < 0.3$  were included in the multinomial model.  
161 Factors shown to be significantly associated with diagnosis included: age ( $P = 0.001$ ), weight



162 ( $P=0.001$ ) and onset of clinical signs ( $P=0.009$ ). These variables were substituted for the  
163 single input variable ‘diagnosis’ in subsequent binary logistic regression.

164

165 Binary logistic regression modelling was performed to identify factors associated with  
166 early postoperative neurological deterioration. Variables were considered for inclusion in  
167 binary logistic regression if  $P<0.30$  and retained in the final model if  $P<0.05$ , based on the  
168 likelihood ratio test. Binary logistic regression was carried out using a Forced Entry Method  
169 to examine associations between included variables with a significance level of  $P<0.05$ .  
170 Results are presented with odds ratios (OR) and 95% confidence intervals (CI) for variables  
171 associated with early postoperative neurological deterioration.

172

## 173 **Results**

174 The study population comprised 100 dogs. Breeds included were Dalmatian ( $n=10$ ),  
175 Labrador Retriever ( $n=9$ ), Great Dane ( $n=8$ ), Dogue de Bordeaux ( $n=7$ ), Bull Mastiff ( $n=7$ ),  
176 Rottweiler ( $n=6$ ), English Cocker Spaniel ( $n=6$ ), Dobermann ( $n=4$ ), Boxer ( $n=3$ ), Basset  
177 Hound ( $n=3$ ), English Pointer ( $n=3$ ), Jack Russell Terrier ( $n=2$ ), Dachshund ( $n=2$ ), West  
178 Highland White Terrier ( $n=2$ ), Beagle ( $n=2$ ) and Bernese Mountain Dog ( $n=2$ ); there were 13  
179 breeds represented by one dog each and 11 crossbreeds. Of the study population 75 dogs  
180 were male and 25 dogs female; overall 45 of the dogs were neutered. At the time of surgery  
181 the median age of dogs was 5.2 years (range 0.4 to 11 years; Table 1). Dogs with OACSM  
182 were significantly younger than dogs with IVDE ( $P=0.001$ ) or meningioma ( $P=0.006$ ). The  
183 median weight was 32.5 kg (range 5 to 80 kg; Table 1). Dogs with OACSM were also  
184 significantly heavier than dogs with IVDE ( $P=0.001$ ), meningioma ( $P=0.004$ ) or vertebral  
185 arch anomalies ( $P=0.024$ ).

186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210

Duration of clinical signs prior to surgery was classified as acute ( $n=15$ ), subacute ( $n=21$ ) or chronic ( $n=64$ ). Dogs underwent a range of advanced imaging diagnostics including myelography ( $n=8$ ), CT-m ( $n=1$ ), MRI ( $n=87$ ) and a combination of CT and MRI ( $n=4$ ). Of those dogs that had myelography or CT-m three had surgery immediately following diagnostics. Diagnoses included OACSM ( $n=41$ ), acute IVDE ( $n=31$ ), meningioma ( $n=11$ ), SAD ( $n=10$ ) and vertebral arch abnormalities ( $n=7$ ; Table 1).

Lesions were located in the cranial cervical region ( $n=27$ ), the caudal cervical region ( $n=59$ ) and affecting both cranial and caudal cervical regions ( $n=14$ ). Seventy dogs had a cervical dorsal laminectomy, 28 had a hemilaminectomy and two dogs had combination of a dorsal laminectomy and hemilaminectomy. A continuous laminectomy was performed in 35 dogs; comprising two sites ( $n=22$ ), three sites ( $n=10$ ) and four sites ( $n=3$ ). Surgical time ranged from 65 to 655 min (median 215 min).

Prior to surgery neurological grade ranged from grade one to five: one ( $n=6$ ), two ( $n=19$ ), three ( $n=46$ ), four ( $n=9$ ) and five ( $n=20$ ). Of the 97 dogs that survived 48h postoperatively, 55% ( $n=52$ ) experienced early postoperative neurological deterioration (Table 1). Postoperative neurological grades ranged from grade one to six: one ( $n=4$ ), two ( $n=6$ ), three ( $n=20$ ), four ( $n=25$ ), five ( $n=41$ ) and six ( $n=1$ ). Of those dogs that demonstrated deterioration in neurological grade, 24 dogs deteriorated by one grade, 23 dogs deteriorated by two grades and five dogs deteriorated by three grades. Of the 52 dogs that experienced early postoperative neurological deterioration 49 dogs had an improved neurological grade at the time of discharge. The remaining three dogs were euthanised during hospitalisation.

211 Other complications included postoperative death within 24 h of surgery ( $n=3$ ),  
212 euthanasia due to neurological deterioration or lack of improvement during hospitalisation  
213 ( $n=3$ ), respiratory compromise requiring postoperative mechanical ventilation ( $n=1$ ), wound  
214 infection requiring antibiotics ( $n=3$ ), severe intraoperative haemorrhage requiring blood  
215 transfusion ( $n=2$ ) and requirement of subsequent surgery at the same site due to incomplete  
216 decompression ( $n=6$ ).

217

218 Duration of hospitalisation ranged from 2 to 28 days (median 7 days; Table 1).  
219 Occurrence of early postoperative neurological deterioration was the only variable  
220 significantly associated with a longer duration of hospitalisation ( $P=0.023$ ).

221

222 Univariate analysis showed that higher bodyweight, younger age, longer duration of  
223 clinical signs, longer surgery times, a higher (more severely affected) neurological grade  
224 before surgery, and diagnosis were significantly associated with early postoperative  
225 neurological deterioration (Table 2). After performing multivariate statistical analysis, four  
226 variables were withheld as independent risk factors for early postoperative neurological  
227 deterioration: diagnosis of OACSM, diagnosis of meningioma, higher (more severely  
228 affected) neurological grade before surgery and longer surgery (Table 3). Diagnoses of  
229 OACSM or meningioma were considered the strongest variables to predict early  
230 postoperative neurological deterioration, followed by higher (more severely affected)  
231 neurological grade and longer surgery time (Table 3).

232

## 233 **Discussion**

234 The results of this study confirm that early postoperative neurological deterioration is  
235 the most frequent complication in dogs undergoing cervical dorsal laminectomy or

236 hemilaminectomy. Identified risk factors for development of this postoperative complication  
237 were diagnosis of OACSM, diagnosis of meningioma, higher (more severely affected)  
238 neurological grade, and longer surgery time.

239

240         The results of this study offer new insights into the controversy surrounding the  
241 morbidity of cervical laminectomies and hemilaminectomies. Previous studies have reported  
242 varying short-term outcomes, complication rates, and hospitalisation times (Gill et al., 1996;  
243 de Risio et al., 2002; da Costa, 2010; Faissler, 2011; De Decker et al., 2012a; Delamide  
244 Gasper et al., 2014). This study indicates that overall 55% of patients undergoing dorsal  
245 cervical decompressive surgeries will experience early postoperative neurological  
246 deterioration, which may be interpreted as a consequence of its invasive nature. Of major  
247 importance is that these results suggest that the specific diagnosis of dogs undergoing these  
248 procedures should be considered the strongest independent variable to predict occurrence of  
249 early postoperative neurological deterioration. Over 75% of patients diagnosed with OACSM  
250 or meningioma experienced postoperative neurological deterioration compared to only 22%  
251 of dogs with acute IVDE. Previous studies reporting high postoperative morbidity with up to  
252 70% of dogs experiencing postoperative deterioration included exclusively patients with  
253 cervical spondylomyelopathy (de Risio et al., 2002). In contrast, those studies that have  
254 suggested more favourable outcomes have predominantly focused on patients diagnosed with  
255 acute IVDE (Gill et al., 1996; Tanaka et al., 2005; Faissler, 2011) or vertebral arch anomalies  
256 (De Decker et al., 2012a). It is therefore possible that the variability in reported postoperative  
257 complications and associated short-term outcomes could largely be explained by the specific  
258 diagnoses of dogs undergoing these procedures.

259

260           It is unclear why patients with OACSM and meningioma are more likely to  
261 experience early postoperative neurological deterioration compared to patients with other  
262 diagnoses. Several authors have suggested that decompression of chronic spinal cord lesions  
263 can lead to reperfusion injuries resulting in acute deterioration after surgery (Rushbridge et  
264 al., 1998; Olby and Jeffery, 2012). However, other disease processes, such as vertebral arch  
265 anomalies and SAD, which are characterised by chronic spinal cord compression, were not  
266 associated with increased risk of early postoperative deterioration. This may suggest that  
267 chronicity of the disease process cannot entirely explain the occurrence of this complication.  
268 Although other reasons cannot be excluded, it is possible that removal of a meningioma is  
269 associated with more spinal cord manipulation and subsequent iatrogenic damage compared  
270 with surgical removal of other causes of spinal cord compression. However, it seems more  
271 difficult to explain the increased risk of early postoperative neurological deterioration in dogs  
272 with OACSM. It has been suggested that dogs with cervical spondylomyelopathy can have a  
273 degree of vertebral instability, which could be aggravated by performing a cervical dorsal  
274 laminectomy (de Risio et al., 2002) and contribute to the development of postoperative  
275 neurological deterioration (de Risio et al., 2002). The role of vertebral instability in the  
276 aetiopathogenesis of cervical spondylomyelopathy is controversial and has not been assessed  
277 objectively (da Costa, 2010; De Decker et al., 2012c).

278

279           Whatever the underlying pathophysiology for early postoperative neurological  
280 deterioration in dogs with OACSM and meningioma, many would consider the occurrence of  
281 this short-term complication in over 75% of patients unacceptable. Although it might be  
282 practically impossible to remove a meningioma without performing a dorsal cervical  
283 laminectomy, alternative treatment modalities can be considered for dogs with OACSM.  
284 Recent studies have evaluated medical management (Delamide et al., 2014) and distraction

285 and stabilization techniques (Lewis et al., 2013) in dogs with OACSM. Both studies  
286 demonstrated promising results, which warrants further investigation into alternative  
287 treatment methods for this disorder. However it should be emphasised that this study  
288 evaluated the occurrence of operative and early postoperative complications in dogs  
289 undergoing dorsal cervical laminectomy or hemilaminectomy and did not evaluate long-term  
290 outcome of dogs undergoing these surgical procedures. This was deemed inappropriate  
291 because of the large variation in included diagnoses, which could have influenced this  
292 variable. It has previously been demonstrated that dogs experiencing early postoperative  
293 neurological deterioration often still experience neurological recovery with good long-term  
294 outcomes (Rushbridge et al., 1998; De Risio et al., 2002) and indeed that was the case with  
295 the dogs in this study. Considering this, it may be appropriate to justify the initial morbidity  
296 associated with cervical dorsal decompressive surgeries for dogs with OACSM and  
297 meningioma. However early postoperative deterioration was found to be significantly related  
298 to prolonged hospitalisation and this may therefore increase costs associated with the  
299 procedure.

300

301         Other identified risk factors for early neurological deterioration after cervical  
302 laminectomy or hemilaminectomy were neurological grade before surgery and duration of  
303 surgery. It is possible that a higher (more severely affected) neurological grade represents  
304 more severe spinal cord pathology and that these dogs are therefore at risk of developing  
305 early postoperative neurological deterioration. Severity of clinical signs in dogs with cervical  
306 spondylomyelopathy is however not necessarily associated with more severe  
307 electrodiagnostic abnormalities, spinal cord compression seen on MRI and worse outcome  
308 after medical treatment (De Decker et al., 2012b). Most sedative and anaesthetic products are  
309 associated with hypotensive effects (Raisis and Musk, 2013). Although special care should be

310 taken to remain adequate spinal cord perfusion during general anaesthesia, it is possible that  
311 prolonged surgeries are associated with an increased risk of spinal hypoperfusion thereby  
312 contributing to a higher risk of early postoperative neurological deterioration (Rossmeisl et  
313 al., 2013). It is also possible that prolonged surgery times are associated with more difficult  
314 surgeries and therefore increased spinal cord manipulation. This hypothesis is however  
315 difficult to prove and remains speculative.

316

317         This study is obviously limited by its retrospective nature. To obtain a large number  
318 of suitable cases, data had to be collected over a number of years and different institutions.  
319 This has potentially limited standardisation of patient assessment and patient care. The  
320 majority of dogs were diagnosed with either OACSM or acute IVDE. This resulted in  
321 unequal group sizes and although this could theoretically have influenced our analysis,  
322 enough patients were included to allow for multivariate statistical analysis. Another potential  
323 limitation is that only a small number of dogs underwent advanced imaging studies after  
324 postoperative neurological deterioration was observed. We can therefore not exclude a  
325 structural cause of early postoperative neurological deterioration in these dogs. However, all  
326 dogs that did not undergo repeat imaging studies demonstrated spontaneous neurological  
327 improvement after variable amounts of time, consistent with previous reports (Rushbridge et  
328 al., 1998; de Risio et al., 2002).

329

### 330 **Conclusions**

331         Patients diagnosed with OACSM and meningioma are significantly more likely to  
332 experience early postoperative neurological deterioration following dorsal cervical  
333 laminectomy and hemilaminectomy than patients undergoing these procedures for the  
334 treatment of other diagnoses. Other risk factors were higher (more severely affected)

335 neurological grade and longer duration of surgery. This information can aid in managing the  
336 expectations of clinical staff and owners with dogs undergoing these procedures.

337

### 338 **Conflict of Interest Statement**

339 None of the authors of this paper has a financial or personal relationship with other  
340 people or organisations that could inappropriately influence or bias the content of this paper.

341

### 342 **Acknowledgments**

343 Preliminary results were presented as an oral abstract at the 27<sup>th</sup> Symposium of the  
344 European College of Veterinary Neurology, Madrid, 18–20 September 2014.

345

### 346 **References**

- 347 da Costa, R.C., 2010. Cervical spondylomyelopathy (wobbler syndrome) in dogs.  
348 *Veterinary Clinics of North America: Small Animal Practice* 40, 881–913.
- 349 De Decker, S., de Risio, L., Lowrie, M., Mauler, D., Beltran, E., Giedja, A., Volk, H.,  
350 2012a. Cervical vertebral stenosis associated with a vertebral arch anomaly in the basset  
351 hound. *Journal of Veterinary Internal Medicine* 26, 1374-1382.
- 352 De Decker, S., Gielen, I., Duchateau, L., Van Soens, I., Oevermann, A., Polis, I., Van  
353 Bree, H., Van Ham, L., 2012b. Evolution of clinical signs and predictors of outcome after  
354 conservative medical treatment for disk-associated cervical spondylomyelopathy in dogs.  
355 *Journal of the American Veterinary Medical Association* 240, 848-857.
- 356 De Decker S., da Costa, R.C., Volk, H., Van Ham, L., 2012c. Current insights and  
357 controversies in the pathogenesis and diagnosis of disk associated cervical  
358 spondylomyelopathy in dogs. *Veterinary Record* 171, 531-537.
- 359 de Risio, L., Munana, K., Murray, M., Olby, N., Sharp, N.J.H., Cuddon, P., 2002.  
360 Dorsal laminectomy for caudal cervical spondylomyelopathy: postoperative recovery and  
361 long-term follow-up in 20 dogs. *Veterinary Surgery* 31, 418-427.
- 362 Delamade Gasper, J.A., Rylander, H., Stenglein, J.L., Waller III, K.R., 2014. Osseous-  
363 associated cervical spondylomyelopathy in dogs: 27 cases (2000-2012). *Journal of the*  
364 *Veterinary Medical Association* 244, 1309-1318.
- 365 Faissler, D., 2011. Cervical Hemilaminectomy: A valid alternative to the ventral slot  
366 procedure. In: *Proceedings of the American College of Veterinary Internal Medicine, Denver,*  
367 *Colorado, 15-18 June 2011, p. 10*
- 368 Gill, P. J., Lippincott, C. L., Anderson, S. M., 1996. Dorsal laminectomy in the  
369 treatment of cervical intervertebral disk disease in small dogs: a retrospective study of 30  
370 cases. *Journal of the American Animal Hospital Association* 32, 77-80.



371 Lewis, M., Olby, N.J., Early, P., 2013. Long-term effect of cervical distraction and  
372 stabilization on neurological status and imaging findings in giant breed dogs with cervical  
373 stenotic myelopathy. *Veterinary Surgery* 42, 701-709.

374 Lipsitz, D., Levitski, R.E., Chauvet, A.E., Berry, W.L., 2001. Magnetic resonance  
375 imaging features of cervical stenotic myelopathy in 21 dogs. *Veterinary Radiology and*  
376 *Ultrasound*, 42, 20–27.

377 Olby, N.J., Jeffey, N.D., 2012. Pathogenesis and Pathophysiology of CNS Disease  
378 and Injury. In: Tobias, K.M., Johnston, S.A. (Eds). *Veterinary Surgery: Small Animal*, First  
379 Ed. Elsevier Saunders, St Louis, Missouri, USA pp 374-387

380 Platt, S.R., da Costa, R.C. 2012. Cervical spine. In Tobias, K.M., Johnston, S.A.  
381 (Eds). *Veterinary Surgery: Small Animal*, First Ed. Elsevier Saunders, St Louis, Missouri,  
382 USA p. 434

383 Rasis, A.L., Musk, G.C., 2013. Analgesia and supportive care. In: Platt, S.R., Olby,  
384 N.J. (Eds). *BSAVA Manual of Canine and Feline Neurology*. Fourth Ed. British Small  
385 Animal Veterinary Association, Gloucester, England, pp 409-432

386 Rossmeis, J.H. Jr, White, C., Pancotto, T.E., Bays, A., Henao-Guerrero, P.N., 2013.  
387 Acute adverse events associated with ventral slot decompression in 546 dogs with cervical  
388 intervertebral disc disease. *Veterinary Surgery*. 42, 795-806

389 Rushbridge, C. Wheeler, S.J., Torrington, A.M., Pead, M.J., Carmichael, S., 1998.  
390 Comparison of two surgical techniques for the management of cervical spondylomyelopathy  
391 in dogs. *Journal of Small Animal Practice* 39, 425-431.

392 Sharp, N.J.H., Wheeler, S.J., 2005. Cervical Spondylomyelopathy. In: *Small animal*  
393 *spinal disorders: diagnosis and surgery*. 2nd ed. St Louis: Elsevier Mosby, pp. 211–232.

394 Tanaka, H., Nakayama, M., Takase, K., 2005. Usefulness of hemilaminectomy for  
395 cervical intervertebral disk disease in small dogs. *Journal of Veterinary Medical Science* 67,  
396 679-683

397

398 **Table 1: Characteristics of dogs undergoing cervical dorsal laminectomy or**  
 399 **hemilaminectomy**

Diagnosis	n	Age Years (Median, Range)	Weight kg (Median, Range)	Neurological grade preoperatively Median (Range)	Neurological grade postoperatively Median (Range) <sup>a</sup>	Neurologically deteriorating <sup>b</sup> n (% , 95% CI)	Days of hospitalisation Median (Range)
OACSM	41	2.7 (0.5-10)	44 (12- 80)	3 (2-4)	5 (0-5)	32 (78%, 65-91)	9 (3-22)
IVDE	31	6.7 (2- 11)	25 (6- 50)	4 (1-5)	4 (1-6)	7 (23%, 7-38)	6 (2-20)
Meningioma	11	7.5 (5- 11)	25 (9- 40)	2 (1-5)	4 (0-5)	9 (82%, 55-109)	6 (4-16)
SAD	10	1 (0.4- 10.3)	28 (5- 59)	3 (3-5)	4 (2-5)	4 (40%, 3-63 )	5 (2-17)
Vertebral arch anomalies	7	1.1 (0.83- 6.6)	32 (22- 48)	3 (2-5)	4 (2-5)	3 (43%, 7-62)	5 (3-12)

400 OACSM, osseous associated cervical spondylomyelopathy; IVDE, intervertebral disc  
 401 extrusion; SAD, spinal arachnoid diverticulum

402 a: assessed at 48 h postoperatively

403 b: those dogs who had deteriorated by one or more neurological grades

404

405 **Table 2: Univariate analysis showing factors associated with early postoperative**  
406 **neurological deterioration**

Factor	P value
Younger age	<i>P</i> =0.030
Higher bodyweight	<i>P</i> =0.039
Higher neurological grade preoperatively (more severely affected)	<i>P</i> =0.001
Dogs having dorsal laminectomy rather than hemilaminectomy	<i>P</i> =0.018
Longer surgery time	<i>P</i> =0.001
Diagnosis	<i>P</i> =0.001

407

408 **Table 3: Results of multivariate analysis showing factors significantly associated with**  
 409 **early postoperative neurological deterioration**  
 410

Factor	P value	Odds ratios	95% Confidence intervals for odds ratios
OACSM diagnosis	<i>P</i> =0.0001	11.4	3.3 to 40.4
Meningioma diagnosis	<i>P</i> =0.014	11.1	1.6 to 75.2
Higher neurological grade preoperatively (more severely affected)	<i>P</i> =0.002	2.3	1.37 to 3.85
Longer surgery time	<i>P</i> =0.014	1.01	1.002 to 1.016

411 OACSM, osseous associated cervical spondylomyelopathy